

**DIRECTIONS FOR USE**

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.  
 CHIPCO® GREEN can be applied year round to:  
 (a) commercial and residential lawns up to 200 m<sup>2</sup>, and patch treatments of up to 5% of estate lawns up to 5,000 m<sup>2</sup> (i.e. 250 m<sup>2</sup>)  
 (b) bowling greens, cricket pitches (excluding outfield), lawn tennis courts and professional sports grounds/stadiums  
 (c) golf greens, tees and patch treatment of up to 5% on fairways.

**GENERAL INFORMATION:**

CHIPCO® GREEN is a contact fungicide containing iprodione which has curative and preventative action against a range of turf diseases. Its special liquid formulation is rainfast within 1 hour and can prevent the build-up of morning dew on grass leaves. 'Switching' to remove this dew is therefore usually unnecessary.

**RESTRICTIONS:**

If grass is being mown, CHIPCO® GREEN should be applied after cutting and further mowing should not take place until AT LEAST 24 hours after treatment.

**APPLICATION RATE:**

For control of Red Thread, Dollar Spot and Fusarium Patch and useful reduction of Anthracnose, Leaf Spot, and Rust, use 20 L/ha of CHIPCO® GREEN in a minimum of 400 - 500 litres of water per hectare of turf.

Application Equipment	Vehicle-mounted Hydraulic Sprayer		Knapsack Sprayer	
	Rate per ha	Water per ha	Rate per 100 m <sup>2</sup>	Water per 100 m <sup>2</sup>
<i>Microdochium nivale</i> (Fusarium Patch), <i>Laetisaria fuciformis</i> (Red Thread) <i>Sclerotinia homeocarpa</i> (Dollar Spot) <i>Colletotrichum cereale</i> (Anthracnose)* <i>Drechslera pose</i> (Leaf Spot)* <i>Puccinia coronata</i> (Rust)*	20 L	500 L	200 ml	5 L

(\*useful reduction when present at time of application against the other listed diseases)

**TIMING OF APPLICATION:**

Curative: Apply at the first sign of disease and repeat as necessary at intervals of 1 month or greater.  
 Preventative: Apply at monthly intervals.  
 CHIPCO® GREEN can be applied year round.

**MIXING AND SPRAYING:**

Add the appropriate quantity of CHIPCO® GREEN to the spray tank, which contains half the required volume of water. Mix thoroughly and make up to the required volume with water. Maintain adequate agitation during application. Continue agitation at all times during spraying and stoppages until the tank is completely empty. Spray immediately after mixing.  
 Wash out all spray equipment thoroughly with water immediately after use.

**Resistance Management:**

CHIPCO® GREEN belongs to the dicarboxamide class of chemistry. There has been no record of managed amenity turf pathogens developing resistance to CHIPCO® GREEN. To avoid the development of resistance, a resistance management strategy should be adopted. This strategy should include using in rotation and/or tank mixture with products that have a different mode of action or different chemical group that are effective against the target disease, as well as non chemical methods of control. Bayer recommends responsible product stewardship to ensure effective long term control of the fungal diseases on this label.

5Le

**CHIPCO® GREEN**  
Contains 25 % w/w Iprodione  
Suspension concentrate (SC)

  
**HARMFUL**

  
**DANGEROUS FOR THE ENVIRONMENT**

Limited evidence of a carcinogenic effect.  
 Very toxic to aquatic organisms, may cause long-term adverse effects in the aquatic environment.

Wear suitable protective clothing and gloves.  
 Keep out of the reach of children.  
 Use appropriate containment to avoid environmental contamination.  
 Keep away from food, drink and animal feeding stuffs.  
 This material and its container must be disposed of in a safe way.  
 If swallowed, seek medical advice and show this container or label.  
 To avoid risks to man and the environment, comply with the instructions for use.

MAPP 13843, PCS 02714



A contact acting fungicide for the control of Fusarium Patch (*Microdochium nivale*), Red Thread (*Laetisaria fuciformis*) and Dollar Spot (*Sclerotinia homeocarpa*) and useful reduction of Anthracnose (*Colletotrichum cereale*), Leaf Spot (*Drechslera pose*) and Rust (*Puccinia coronata*) where present at time of application, on managed amenity turf.\* (\*See Important Information)

**Fungicide**

F

**SAFETY PRECAUTIONS**

**Operator Protection**  
 TAKE OFF IMMEDIATELY heavily contaminated clothing.  
 WASH SPLASHES OF CONCENTRATE from skin and eyes immediately.  
 When using do not eat drink or smoke.  
 WASH HANDS before meals and after work.  
 In case of accident or if you feel unwell seek medical advice immediately (show the label where possible).

**Environmental Protection**

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing water body, unless a Local Environment Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 m of the top of a ditch which is dry at the time of application. DO NOT ALLOW DIRECT SPRAY from hand-held sprayers to fall within 1 m of the top of the bank of a static or flowing water body. Aim spray away from water. – UK only.

**Storage and Disposal**

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.  
 WASH OUT CONTAINER THOROUGHLY and dispose of safely.  
 DO NOT RE-USE CONTAINER for any purpose.

PROTECT FROM FROST.  
 SHAKE THE CONTAINER WELL BEFORE USE.

WHO-classification: III (Slightly hazardous).

**Important Information:**

FOR USE ONLY AS AN AMENITY/ HORTICULTURAL FUNGICIDE RESTRICTED TO PROFESSIONAL USERS.  
 For use on managed amenity turf (see 'Other Specific Restrictions').  
 Maximum individual dose: 20 L/ha.  
 Maximum number of treatments: 4 per year.  
 Other Specific Restrictions:  
 A minimum of four weeks must be observed between treatments.  
 For use on:  
 (a) commercial and residential lawns up to 200 m<sup>2</sup>, and patch treatments of up to 5% of estate lawns up to 5,000 m<sup>2</sup> (i.e. 250 m<sup>2</sup>)  
 (b) bowling greens, cricket pitches (excluding outfield), lawn tennis courts and professional sports grounds/stadiums  
 (c) golf greens, tees and patch treatment of up to 5% on fairways.

This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with PSD's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years. – UK only.

LERAP  
B


**Read the label before use. Using this product in a manner that is inconsistent with the label may be an offence.**  
**Follow the code of practice for using plant protection products.**

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work - UK only.

**For detailed directions for use please see inside leaflet.**

Chipco Green - GB - 5 L - GB79444060E - ARTICLE 04203959

For 24 hour emergency information contact Bayer CropScience Limited.  
 Telephone: 0800 220876 or nearest National Poisons Information Centre.  
 +44(0)1603-242424 (Overseas 24 hr)

 This label is compliant with the CPA Voluntary Initiative Guidance - UK only.

Batch number : See on the pack

Authorization holder:  
**Bayer S.A.S.**  
*Bayer Environmental Science*  
 16 rue Jean-Marie Leclair - CS 90106  
 69266 LYON CEDEX 09 - FRANCE

Bayer Environmental Science  
 a Business Group of Bayer CropScience Limited  
 230 Cambridge Science Park  
 Milton Road  
 Cambridge CB4 0WB  
 Tel : 00800 1214 9451  
 www.escience.bayercropscience.co.uk

