

## Holster XL

Holster XL is a turf herbicide that has been specifically developed for the control of problem weeds in amenity turf (including fine turf). Weeds treated with Holster XL show symptoms such as twisted leaf expansion and altered growth - ultimately leading to the decay of the plant.

Holster provides outstanding control of a range of broad-leaved weeds including Suckling Clover, Slender Speedwell, Daisy and Common Mouse Ear.

Apply when the weeds are actively growing (normally between April and September) and when the soil is moist. Best results will be obtained by application in spring or early summer before weeds begin to flower. New turf may be treated in the spring provided that at least two months have elapsed between sowing and treatment.

### Product Advantages:

- Can be applied over new grass only two months after sowing
- Excellent grass safety
- Low application rates - 2 litres per hectare for cost effective control
- Contains three powerful active ingredients
- Controls many hard to kill weeds in turf

### Specifications:

<b>Pack Size:</b>	<b>MAPP No.</b> 16407
2 ltr	<b>LERAP:</b> Category n/a
<b>Type:</b>	<b>Contains:</b> 105 g/ltr Fluroxypyr,
Liquid	285 g/ltr 2,4-D &
<b>Pack Coverage:</b>	52.5 g/ltr Dicamba
10,000 sq. m	



Available in stock from Collier Turf Care now  
Call your local Technical Sales Rep to place your order

## Holster XL

### Rates of Use:

Method of Application	Rate of Holster	Water Rate	Area Treated
Powered Sprayer	2 litres	200 - 500 litres	1 hectare
Knapsack Sprayer	20 ml	5 - 10 litres	100 sq. m
Turfguard Professional	100 ml	25 litres	500 sq. m

### Weed Control:

Autumn Hawkbit	Dandelion	Self Heal
Black Medick	Docks	Silverweed
Bulbous Buttercup	Heath Bedstraw	Sorrel
Cat's-Ear	Field Madder	Sea Milkwort
Charlock	Knotgrass	Slender Speedwell
Cinquefoil	Mayweeds	Thistles
Cleavers	Mouse-ear Hawkweed	Trefoils
Common Stork's-bill	Meadow Buttercup	Thrift
Common Chickweed	Parsley Piert	White Clover
Creeping Buttercup	Procumbent Pearlwort	Yarrow
Common Mouse-ear	Plantains	Yellow Rattle
Daisy	Common Ragwort	Yellow Suckling Clover

Do not apply if the turf is wet or is rainfall is expected within 4 hours of application.  
Do not spray in periods of drought (unless irrigation is applied)  
Use an appropriate water volume to achieve good coverage of the weeds. Apply as a medium spray as defined by the British Crop Protection Council.



Available in stock from Collier Turf Care now  
Call your local Technical Sales Rep to place your order