

Enforcer[®]

Selective herbicide
for the control of
broad-leaved
weeds in managed
amenity turf

Where needs take us

AICL

Enforcer®

Enforcer is a selective herbicide with foliar and some root activity for use against a range of common broad-leaved weeds such as White Clover (*Trifolium repens*), Creeping Buttercup (*Ranunculus repens*), Daisy (*Bellis perenis*) and Dandelion (*Taraxacum officinale*).

Key Points

- Potent combination of four active ingredients
- Ideal for all sports turf areas and lawns
- Various application methods including knapsack



Directions for use

1. Shake well before use.
2. Enforcer can be applied by tractor-mounted boom sprayer or by knapsack sprayer.
3. The activity of Enforcer is promoted by climatic conditions which favour photosynthesis such as warm temperatures and sunny conditions.
4. Avoid treatment during dry periods and when rain is expected within 24 hours after the treatment.
5. Do not spray to grass suffering from drought or during freezing conditions.
6. May take up to 3 weeks before weed destruction is seen. Do not re-apply if weed destruction is not seen directly after treatment.
7. For best results, apply the product at least 5 days after the last mowing and wait 7 days after application before mowing again.
8. Do not use on new grass for 6 months after establishment.
9. When renovating treated areas an interval of 6-8 weeks should elapse before re-seeding.

Recommended period of use

April - September.

Product characteristics

Pack size:	5 litre bottle
Packs per outer:	4 x 5 litre
Contains:	70 g/L 2,4-D, 20 g/L dicamba, 70 g/L MCPA and 42 g/L mecoprop-P MAPP no. 17274 Not available in ROI

Health and safety

A Material Safety Data sheet is available on request.

Application rate

Maximum individual dose:	7.5 L/ha
Maximum number of treatments:	1 per year
Latest time of application:	End of September in the year of treatment
Water Volume:	1,000 L/ha
Maximum total dose:	7.5 L/ha

Weeds Controlled

Weed	Application rate	Comments
White clover (<i>Trifolium repens</i>), daisy (<i>Bellis perenis</i>), deadnettle (<i>Lamium sp.</i>), speedwell (<i>Veronica sp.</i>), Shepherd's Purse (<i>Capsella bursa pastoris</i>) and Field Forget-me-not (<i>Myosotis arvensis</i>)	7.5 L/ha	Susceptible
Dandelion (<i>Taraxacum officinale</i>), broad-leaved dock (<i>Rumex obtusifolius</i>), Self heal (<i>Prunella vulgaris</i>), creeping buttercup (<i>Ranunculus repens</i>), yarrow (<i>Achillea millefolium</i>) and corn chamomile (<i>Anthemis arvensis</i>).	7.5 L/ha	Moderately susceptible