



# TROPHY XL®

## New turf herbicide for challenging weed control



# TROPHY XL<sup>®</sup>



Trophy XL<sup>®</sup> combines a new active ingredient, Arylex™, with established herbicide technology, offering broad spectrum weed control in turf. Its advanced formulation acts quickly on actively growing weeds and is highly effective against persistent weeds like plantain and clover.

Trophy XL is an EC (Emulsifiable Concentrate) containing 280 g/L fluroxypyr, 12.5 g/L halauxifen-methyl (marketed as Arylex), and 12 g/L cloquintocet-mexyl as a safener. This post-emergence herbicide is designed for use on managed amenity turf, targeting a wide range of broad-leaved weeds.

## Key Points

- **Broad-spectrum, selective herbicide for extensive weed control**
- **Rapid and effective control of Greater plantain and Common plantain**
- **Robust control of actively growing weeds in cooler weather**
- **Low-odour formulation for a more pleasant application experience**
- **Can be applied in a single treatment annually**
- **Suitable for newly established or mature turf areas**
- **Designed to integrate smoothly with existing turf maintenance schedule**

## Product characteristics

Formulation type:	Liquid EC (Emulsifiable Concentrate)
Technology:	Selective herbicide
Pack size:	1 litre
Pack coverage:	1.67 hectares
MAPP:	20820
Contains:	80 g/L fluroxypyr, 12.5 g/L halauxifen-methyl (marketed as Arylex), and 12 g/L cloquintocet-mexyl (safener)

## Application information

Maximum individual dose	0.6 L/ha
Application rate:	0.6 L/ha
Water volume:	200–500 L/ha
Maximum number of treatments:	1 per year
Use period:	1st April to 30th September
Application area:	Managed amenity turf

# Effective weed management programmes with Trophy XL® & Praxys®

Trophy XL and Praxys are selective herbicides designed to control a broad spectrum of weeds. The programmes below outline different approaches for using each product, depending on weed types and infestation levels.



## Single treatment programmes

Trophy XL and Praxys provides strong, targeted control of broad-leaved weeds with a single treatment.

### Trophy XL

Ideal for sites with a range of broad-leaved weeds, especially effective on plantain.

**Application timing:**  
Apply Trophy XL once in April, May, or June.

### Praxys

Suitable for areas with a variety of broad-leaved weeds, particularly effective on daisy.

**Application timing:**  
Apply Praxys once in April, May, or June.

## Two treatment programme

For areas prone to reinfestation and where multiple broad-leaved weeds persist, a two-treatment approach is recommended to maintain control throughout the season.

### 1st Treatment

Apply Praxys in April or May for initial control of broad-leaved weeds and daisy.

### 2nd Treatment

Apply Trophy XL in June, July, or August for continued suppression and to address weeds like plantain and emerging infestations.

**This sequential treatment provides ongoing coverage and mitigates the resurgence of common broad-leaved weeds.**

## Flexible application window

Both Trophy XL and Praxys can be applied any time between April and September, depending on conditions and weed pressure. Regularly inspect turf and treat when weeds are present and actively growing for optimal results.

## Encouraging grass competitiveness

As part of an integrated weed management plan adequate nutrition is essential for sustaining turf health.

Always read the label and product information before use.



Epsilon House, West Road,  
Ipswich, Suffolk IP3 9FJ  
T: +44 (0)1473 237100  
E: [prof.sales@icl-group.com](mailto:prof.sales@icl-group.com)



[www.icl-growingsolutions.uk](http://www.icl-growingsolutions.uk)



Trophy XL, Praxys and Arylex are  
Trademarks of Corteva Agriscience  
and its affiliate companies.