

WEDB in annihol yranstand. Het Dis yn annihol yranstand. He (COSHR) Control of Substances Haardous to Health Replations my adoly to the use of this product a use (Ukon RED DIRECTIONS FOR USE on ATTACHED LEAFLET. TRIVER FROM PROST. TRIVER RIVES CONTAINERS, NUNCTURE AND INVERT TO DRY A THE OF USE Dis Aprodicinces Limited Prix Capoli Invest.

Mees Park, Fulbourn, Cumbridge C821 SXE.
 Stephone: 01462 4267272 Fax:01462 426605
 St How Emergency TelephoneNumber: 01553 761251
 Technical Hotline: 0800 689 689, e-mailtukhofline@control

NET CONTENTS : 1 LITRE

Nothing **gets to grips with weeds** like Icade



icade

For the control of invasive and tough woody weeds


Icade quickly translocates throughout the target weed



Areas where lcade can be used include

Railway and motorway embankments, industrial areas, under powerlines, cemeteries, utilities, parkland, around greenkeeper sheds, golf course roughs and areas of grassland of no agricultural interest. Icade herbicide is approved for use in Amenity Grassland. Icade is NOT approved for use on airfields. Do not use treated area for animal feed or bedding.



























Get the best out of lcade



- 🔗 Apply Icade to actively growing weeds with a dry leaf
- Spray Icade to thoroughly wet the leaves up to the point of run off
- Sensure good spray coverage of the whole weed
- 🗸 Spray to avoid drift
- 🧭 Allow at least 1 hour before any rainfall
- Only mix enough spray for the job in hand

Application information

Knapsack application only

Apply at 150ml in 10L of water for the control of Bramble and Gorse. Apply at 200 ml in 200L of water for the control of Japanese Knotweed, Buddleia and Broom. Do not spray up to drop line of trees. 5 metre buffer zone to water.

Weed Control

Weeds controlled, rates of use and timing of application.

Weeds	Rates of use ml/10 litres of water	Optimum timing of application
Bramble	150	up to 1 metre high
Broom	150	up to 1 metre high
Buddleia	150	up to 1 metre high
Gorse	150	up to 1 metre high
Japanese Knotweed	150	up to 1 metre high, with good folliage cover

When to use Icade

For best results apply lcade early in the season during active growth. For annual applications do not use between 31st July and 1st March.

- For applications made between 1st March and 1st June,
 1 application may be made per calendar year
- For applications made between 1st June and 1st August, only 1 application may be made in a 2 year period
 - For applications made between 1st August and 31st August, only 1 application may be made in a 3 year period.

	Jan	Feb	Mar	Apr	Мау	Jun	Jul	Aug	Sep	Oct	Nov	Dec
--	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----



Why use Icade?

Unwanted vegetation in amenity grassland and non crop areas causes safety hazards, severe economic loss, and damage to desirable structures. As well as being unsightly, invasive weeds also reduce biodiversity and can pose serious risk to human health.

With the need for the most efficient weed killing technology, whilst being safe to humans and the environment, Corteva has combined the latest technology in Icade.

Key points

V	Effective against a wide range of tough and invasive weeds – including Japanese Knotweed and Buddleia
Ø	Handy 1 litre pack size
V	No hazard classification - favourable COSHH assessment
Ø	Virtually no odour
	Rainfast in 1 hour
	Safe to grasses
V	Outstanding levels of translocation gets ICADE deep into roots of the weeds



USE HERBICIDES SAFELY. ALWAYS READ THE LABEL AND PRODUCT INFORMATION BEFORE USE. Corteva Agriscience, CPC2 Capital Park, Fulbourn, Cambridge, CB21 5XE Tel: (01462) 457272 Fax: (01462) 426605 24 Hour Emergency Telephone Number: +44 (0) 161 884 1235 e-mail: ukhotline@corteva.com Technical Hotline: 0800 689 8899

Icade is a trademark of Dow AgroSciences LLC. Icade contains 12 g/litre aminopyralid and 120 g/litre of triclopyr. MAPP No. 19069

Epsilon House West Road Ipswich Suffolk IP3 9FJ United Kingdom

+44 1473 237 123 prof.sales@icl-group.com

www.icl-sf.co.uk www.icl-sf.ie

